

# Algorithms and bounds for hidden sums in cryptographic trapdoors

C. Brunetta<sup>a</sup>, M. Calderini<sup>a</sup>, M. Sala<sup>a</sup>

<sup>a</sup>*Department of Mathematics, University of Trento, Via Sommarive 14, 38100 Povo (Trento), Italy*

---

## Abstract

Sometimes it is possible to embed an algebraic trapdoor in a block cipher. Building on previous research, in this paper we investigate an especially dangerous algebraic structure, which is called a hidden sum and which is related to some regular subgroups of the affine group. Mixing group theory arguments and cryptographic tools, we pass from characterizing our hidden sums to designing an efficient algorithm to perform the necessary preprocessing for the exploitation of the trapdoor.

*Keywords:* Hidden sums, trapdoor, mixing layer, cryptography, block ciphers.

---

## 1. Introduction

Sometimes it is possible to embed an algebraic trapdoor in a block cipher. Building on previous research, in this paper we investigate an algebraic structure, which is called a hidden sum and which is related to some regular subgroups of the affine group. To be more precise, in [1], the authors study some elementary abelian regular subgroups of the affine general linear group acting on a space  $V = (\mathbb{F}_2)^N$ , in order to construct a trapdoor for a class of block ciphers. These subgroups induce alternative operations  $\circ$  on  $V$ , so that  $(V, \circ)$  results a vector space over  $\mathbb{F}_2$ . In [1], it is shown that for a class of this operations, called *hidden sums*, it is possible to represent the elements with respect to a fixed basis of  $(V, \circ)$  in polynomial time. Moreover, an estimate on the number of these operations is given. Using this class of hidden sums, the authors provide an attack, which works in polynomial time, on ciphers that are vulnerable to this kind of trapdoor.

In this article we continue the analysis started in [1]. In Section 3 we

---

*Email addresses:* `brunetta@unitn.it` (C. Brunetta), `marco.calderini@unitn.it` (M. Calderini), `maxsalacodes@gmail.com` (M. Sala)

give a lower bound on the number of hidden sums, comparing also this value with a previous upper bound. In Section 4 we study the problem of determining the possible maps that are linear with respect to these hidden sums. More precisely, we provide an algorithm that takes in input a given linear map (with respect to the usual XOR on  $V$ ) and returns the hidden sums for which this map is linear also with respect to these. Our aim is to individuate a family of hidden sums that can weaken the components of a given cipher, or to design a cipher containing the trapdoor based on hidden sums. In the last section we apply the procedure given in Section 3 to the mixing layer of the cipher PRESENT [2], obtaining a set of hidden sums which linearize this permutation matrix and might in principle be used to attack the cipher.

## 2. Preliminaries and motivations

For any positive integer  $m$ , we let  $[m] = \{1, \dots, m\}$ . We write  $\mathbb{F}_q$  to denote the finite field of  $q$  elements, where  $q$  is a power of prime, and  $(\mathbb{F}_q)^{s \times t}$  to denote the set of all matrices with entries over  $\mathbb{F}_q$  with  $s$  rows and  $t$  columns. The identity matrix of size  $s$  is denoted by  $I_s$ . We use

$$\mathbf{e}_i = (\underbrace{0, \dots, 0}_{i-1}, 1, \underbrace{0, \dots, 0}_{N-i}) \in (\mathbb{F}_q)^N$$

to denote the unit vector, which has a 1 in the  $i$ th position, and zeros elsewhere. Let  $m \geq 1$ , the vector (sub)space generated by the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is denoted by  $\text{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ .

Let  $V = (\mathbb{F}_q)^N$ , we denote by  $\text{Sym}(V)$ ,  $\text{Alt}(V)$ , respectively, the symmetric and the alternating group acting on  $V$ . By  $\text{AGL}(V)$  and  $\text{GL}(V)$  we denote the affine and linear group of  $V$ . We write  $\langle g_1, \dots, g_m \rangle$  for the group generated by  $g_1, \dots, g_m$  in  $\text{Sym}(V)$ . The map  $1_V$  will denote the identity map on  $V$ .

Let  $G$  be a finite group acting on  $V$ . We write the action of a permutation  $g \in G$  on a vector  $\mathbf{v} \in V$  as  $\mathbf{v}g$ .

We denote the translation with respect to a vector  $\mathbf{v} \in V$  by  $\sigma_{\mathbf{v}} : \mathbf{x} \mapsto \mathbf{x} + \mathbf{v}$ .

### 2.1. Block ciphers and hidden sums

Most modern block ciphers are iterated ciphers, i.e. they are obtained by the composition of a finite number  $\ell$  of rounds.

Several independent formal definitions have been proposed for an iterated block cipher, e.g. *substitution permutation network* [3] and *key-alternating block cipher* [4]. Here we present one more recent definition [5] that determines a class large enough to include some common ciphers (AES [6], SERPENT [7], PRESENT [2]), but with enough algebraic structure to allow for security proofs.

Let  $V = (\mathbb{F}_2)^N$  with  $N = mb$ ,  $b \geq 2$ . The vector space  $V$  is a direct sum

$$V = V_1 \oplus \cdots \oplus V_b,$$

where each  $V_i$  has the same dimension  $m$  (over  $\mathbb{F}_2$ ). For any  $\mathbf{v} \in V$ , we will write  $\mathbf{v} = \mathbf{v}_1 \oplus \cdots \oplus \mathbf{v}_b$ , where  $\mathbf{v}_i \in V_i$ .

Any  $\gamma \in \text{Sym}(V)$  that acts as  $\mathbf{v}\gamma = \mathbf{v}_1\gamma_1 \oplus \cdots \oplus \mathbf{v}_b\gamma_b$ , for some  $\gamma_i$ 's in  $\text{Sym}(V_i)$ , is a *bricklayer transformation* (a “parallel map”) and any  $\gamma_i$  is a *brick*. Traditionally, the maps  $\gamma_i$ 's are called S-boxes and  $\gamma$  a “parallel S-box”. A linear map  $\lambda : V \rightarrow V$  is traditionally said a “Mixing Layer” when used in composition with parallel maps. For any  $I \subset [b]$ , with  $I \neq \emptyset, [b]$ , we say that  $\bigoplus_{i \in I} V_i$  is a *wall*.

**Definition 2.1.** A linear map  $\lambda \in \text{GL}(V)$  is a proper mixing layer if no wall is invariant under  $\lambda$ .

We can characterize the translation-based class by the following:

**Definition 2.2.** A block cipher  $\mathcal{C} = \{\varphi_k \mid k \in \mathcal{K}\} \subset \text{Sym}(V)$ , where  $\mathcal{K}$  is the set containing all the session keys, over  $\mathbb{F}_2$  is called translation based (tb) if:

- it is the composition of a finite number of rounds, such that any round  $\rho_{k,h}$  can be written<sup>1</sup> as  $\gamma\lambda\sigma_{\bar{k}}$ , where
  - $\gamma$  is a round-dependent bricklayer transformation (but it does not depend on  $k$ ),
  - $\lambda$  is a round-dependent linear map (but it does not depend on  $k$ ),
  - $\bar{k}$  is in  $V$  and depends on both  $k$  and the round ( $\bar{k}$  is called a “round key”),
- for at least one round, which we call proper, we have (at the same time) that  $\lambda$  is proper and that the map  $\mathcal{K} \rightarrow V$  given by  $k \mapsto \bar{k}$  is surjective.

For a tb cipher it is possible to define the following groups. For each round  $h$

$$\Gamma_h(\mathcal{C}) = \langle \rho_{k,h} \mid k \in \mathcal{K} \rangle \subseteq \text{Sym}(V),$$

and the round function group is given by

$$\Gamma_\infty(\mathcal{C}) = \langle \Gamma_h(\mathcal{C}) \mid h = 1, \dots, \ell \rangle.$$

An interesting problem is determining the properties of the permutation group  $\Gamma_\infty(\mathcal{C}) = \Gamma_\infty$  that imply weaknesses of the cipher. A *trapdoor* is a

---

<sup>1</sup>we drop the round indices

hidden structure of the cipher, whose knowledge allows an attacker to obtain information on the key or to decrypt certain ciphertexts.

The first paper dealing with properties of  $\Gamma_\infty$  was published by Paterson [8], who showed that if this group is imprimitive, then it is possible to embed a trapdoor in the cipher. On the other hand, if the group is primitive no such trapdoor can be inserted.

However, the primitivity of  $\Gamma_\infty$  does not guarantee the absence of trapdoors. Indeed, if the group is contained in  $\text{AGL}(V)$ , the encryption function is affine and once we know the image of a basis of  $V$  and the image of the zero vector, then we are able to reconstruct the matrix and the translation that compose the map. This motivated the authors of [1] on studying different abelian operations, which can be individuated on  $V$ , called *hidden sums*, in order to embed a trapdoor in a block cipher.

In the remainder of this section, we summarize the theory presented in [1, Section 2-3-5].

**Remark 2.3.** *If  $T$  is an elementary abelian regular group, there exists a vector space structure  $(V, \circ)$  such that  $T$  is the related translation group. In fact, being  $T$  regular the elements of the group can be labelled*

$$T = \{\tau_{\mathbf{a}} \mid \mathbf{a} \in V\},$$

where  $\tau_{\mathbf{a}}$  is the unique map in  $T$  such that  $0 \mapsto \mathbf{a}$ . Then, the sum between two elements is defined by  $\mathbf{x} \circ \mathbf{a} := \mathbf{x} \tau_{\mathbf{a}}$ . Clearly,  $(V, \circ)$  is an abelian additive group and thus a vector space over  $\mathbb{F}_2$ .

Following [1] a hidden sum on  $V$  is an abelian operation  $\circ$  such that  $(V, \circ)$  is a vector space.

In the following, with the symbol  $+$  we refer to the usual sum over the vector space  $V$ . We denote by  $T_+ = T(V, +)$ ,  $\text{AGL}(V, +)$  and  $\text{GL}(V, +)$ , respectively, the translation, affine and linear groups w.r.t.  $+$ . We use  $T_\circ$ ,  $\text{AGL}(V, \circ)$  and  $\text{GL}(V, \circ)$  to denote, respectively, the translation, affine and linear groups corresponding to a hidden sum  $\circ$ .

From [1] we have three interesting problems:

1. Determine the operations  $\circ$  (equivalently the translation groups  $T_\circ$ ) such that  $T_+ \subseteq \text{AGL}(V, \circ)$ .
2. Determine the operations  $\circ$  such that  $T_\circ \subseteq \text{AGL}(V, +)$ .
3. Given a parallel S-box  $\gamma$  and a mixing layer  $\lambda$ , determine the operations  $\circ$  such that  $\gamma, \lambda \in \text{AGL}(V, \circ)$  or  $\gamma\lambda \in \text{AGL}(V, \circ)$ .

The following vector space plays an important role for studying these problems. Let  $T$  be any subgroup of the affine group, we can define the vector space

$$U(T) = \{\mathbf{v} \in V \mid \sigma_{\mathbf{v}} \in T\}.$$

In [1] the authors prove

**Proposition 2.4** (Proposition 3.6 [1]). *Let  $V = (\mathbb{F}_2)^N$ . Let  $T \subseteq \text{AGL}(V, +)$  be an elementary abelian regular subgroup. If  $T \neq T_+$ , then  $1 \leq \dim(U(T)) \leq N - 2$ .*

A characterization is given in [1] for the maps that generate a translation group  $T_\circ \subseteq \text{AGL}(V, +)$  such that  $T_+ \subseteq \text{AGL}(V, \circ)$ . We recall that for every  $\mathbf{a}$ ,  $\tau_{\mathbf{a}} \in T_\circ \subseteq \text{AGL}(V, +)$  can be written as  $\kappa_{\mathbf{a}}\sigma_{\mathbf{a}}$  for a linear map  $\kappa_{\mathbf{a}} \in \text{GL}(V, +)$ . We will denote by  $\Omega(T_\circ) = \{\kappa_{\mathbf{a}} \mid \mathbf{a} \in V\} \subset \text{GL}(V, +)$ . Moreover  $\kappa_{\mathbf{a}} = 1_V$  if and only if  $\mathbf{a} \in U(T_\circ)$

**Lemma 2.5** (Lemma 3.4 [1]). *Let  $T_\circ \subseteq \text{AGL}(V, +)$ . We have that  $\Omega(T_\circ)$  is unipotent. Moreover, for all  $\mathbf{y} \in V$   $\kappa_{\mathbf{y}}^2 = 1_V$ .*

**Theorem 2.6** (Theorem 3.17 [1]). *Let  $V = (\mathbb{F}_2)^{n+d}$ , with  $n \geq 2$ ,  $d \geq 1$ , and  $T_\circ \subseteq \text{AGL}(V, +)$  be such that  $U(T_\circ) = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$ . Then,  $T_+ \subseteq \text{AGL}(V, \circ)$  if and only if for all  $\kappa_{\mathbf{y}} \in \Omega(T_\circ)$  there exists a matrix  $B_{\mathbf{y}} \in (\mathbb{F}_2)^{n \times d}$  such that*

$$\kappa_{\mathbf{y}} = \begin{bmatrix} I_n & B_{\mathbf{y}} \\ 0 & I_d \end{bmatrix}.$$

Note that we can always suppose that  $U(T_\circ)$  is generated by the last vectors of the canonical basis, as any group  $T_\circ$  is conjugated to a group  $T_{\circ'}$  such that  $U(T_{\circ'}) = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$  (see [1, Theorem 3.14]).

**Remark 2.7.** *Since  $U(T_\circ)$  is generated by the last vectors of the canonical basis then the maps  $\tau_{\mathbf{e}_i}$  generate  $T_\circ$ , i.e. the canonical vectors form a basis also for the vector space  $(V, \circ)$ .*

In [1] it turns out that for any hidden sum  $T_\circ \subseteq \text{AGL}(V, +)$ , any given vector  $\mathbf{v}$  can be represented in  $(V, \circ)$ , with respect to the canonical vectors, in polynomial time ( $\mathcal{O}(N^3)$ ). From this the authors prove that a hidden sum trapdoor is practical, whenever  $\Gamma_\infty \subseteq \text{AGL}(V, \circ)$  and  $T_\circ \subseteq \text{AGL}(V, +)$ . A class of these hidden sums is used in [9] to weaken the non-linearity of some APN S-boxes. In [9], also a differential attack with respect to hidden sums is presented.

An upper bound on the number of some hidden sums is given in [1] and reported below.

Let  $n \geq 2$  and  $d \geq 1$  and define for all  $1 \leq i \leq n$  the matrix

$$\kappa_{\mathbf{e}_i} = \begin{bmatrix} & b_{1,1}^{(i)} & \dots & b_{1,d}^{(i)} \\ I_n & \vdots & & \vdots \\ & b_{n,1}^{(i)} & \dots & b_{n,d}^{(i)} \\ 0 & & I_d & \end{bmatrix}. \quad (1)$$

**Theorem 2.8** (Theorem 5.5 [1]). *Let  $N = n + d$  and  $V = (\mathbb{F}_2)^N$ , with  $n \geq 2$*

and  $d \geq 1$ . The elementary abelian regular subgroups  $T_o \subseteq \text{AGL}(V, +)$  such that  $\dim(U(T_o)) = d$  and  $T_+ \subseteq \text{AGL}(V, \circ)$  are

$$\begin{bmatrix} N \\ d \end{bmatrix}_2 \cdot |\mathcal{V}(\mathcal{I}_{n,d})| \quad (2)$$

where  $\mathcal{V}(\mathcal{I}_{n,d})$  is the variety of the ideal  $\mathcal{I}_{n,d} \subset \mathbb{F}_2[b_{i,j}^{(s)} \mid i, s \in [n], j \in [d]]$  generated by

$$\mathcal{S}_0 \cup \mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_3$$

with

$$\begin{aligned} \mathcal{S}_0 &= \left\{ \left( b_{i,j}^{(s)} \right)^2 - b_{i,j}^{(s)} \mid i, s \in [n], j \in [d] \right\} \\ \mathcal{S}_1 &= \left\{ \prod_{i=1}^n \prod_{j=1}^d \left( 1 + \sum_{s \in S} b_{i,j}^{(s)} \right) \mid S \subseteq [n], S \neq \emptyset \right\}, \\ \mathcal{S}_2 &= \left\{ b_{i,j}^{(s)} - b_{s,j}^{(i)} \mid i, s \in [n], j \in [d] \right\}, \\ \mathcal{S}_3 &= \left\{ b_{i,j}^{(i)} \mid i \in [n], j \in [d] \right\}, \end{aligned}$$

and  $\begin{bmatrix} N \\ d \end{bmatrix}_q = \prod_{i=0}^{d-1} \frac{q^{N-i}-1}{q^{d-i}-1}$  is the Gaussian Binomial.

**Proposition 2.9** (Proposition 5.6 [1]). *Let  $\mathcal{I}_{n,d}$  defined as in Theorem 2.8, then*

$$|\mathcal{V}(\mathcal{I}_{n,d})| \leq 2^{d \frac{n(n-1)}{2}} - 1 - \sum_{r=1}^{n-2} \binom{n}{r} \prod_1^{n-r} (2^d - 1) = \mu(n, d).$$

### 3. New lower bounds and asymptotic estimates

In this section we will provide a lower bound on the cardinality of the variety  $\mathcal{V}(\mathcal{I}_{n,d})$ . Moreover we will show that the ratio between the upper bound and lower bound is less than  $e^{\frac{2^d+1}{2^d(2^d-1)}}$ .

From Theorem 2.6 and from Remark 2.7, we have that a group  $T_o$  with  $U(T_o) = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$  is determined by the maps  $\kappa_{\mathbf{e}_i}$ 's, and in particular by  $B_{\mathbf{e}_1}, \dots, B_{\mathbf{e}_n}$  ( $B_{\mathbf{e}_i} = 0$  for  $n+1 \leq i \leq n+d$ ). From Theorem 2.8 we have a one-to-one correspondence between the matrices  $B_{\mathbf{e}_i}$ 's and the points of  $\mathcal{V}(\mathcal{I}_{n,d})$ . In particular, we need to construct the matrices  $B_{\mathbf{e}_1}, \dots, B_{\mathbf{e}_n}$  so that:

- i)  $\dim(U(T_o)) = d$ . Then for any  $\mathbf{v} \in \langle \mathbf{e}_1, \dots, \mathbf{e}_n \rangle$ ,  $B_{\mathbf{v}} \neq 0$ . Indeed,  $\mathbf{v} \in U(T_o)$  if and only if  $B_{\mathbf{v}} = 0$ . Moreover, for any  $\mathbf{v} = (v_1, \dots, v_{n+d}) \in V$  we have

$$\kappa_{\mathbf{v}} = \begin{bmatrix} I_n & \sum_{i=1}^n v_i B_{\mathbf{e}_i} \\ 0 & I_d \end{bmatrix}$$

and so we get that  $B_{\mathbf{v}} = \sum_{i=1}^n v_i B_{\mathbf{e}_i}$  (see Lemma 5.4 in [1]).

This implies that every non-null linear combination of  $B_{\mathbf{e}_1}, \dots, B_{\mathbf{e}_n}$  is non-zero. Note that, in Theorem 2.8, this condition is given by  $\mathcal{S}_1$ .

- ii)  $T_{\circ}$  is abelian. This happens if and only if the  $i$ th row of  $B_{\mathbf{e}_j}$  is equal to the  $j$ th row of  $B_{\mathbf{e}_i}$ , obtaining the set  $\mathcal{S}_2$ .
- iii)  $T_{\circ}$  is elementary, that is,  $\tau_{\mathbf{v}}^2 = 1_V$  for all  $\mathbf{v}$ . Then  $\kappa_{\mathbf{v}}$  fixes  $\mathbf{v}$  and, in particular,  $\kappa_{\mathbf{e}_i}$  fixes  $\mathbf{e}_i$ . This is equivalent to having the  $i$ th row of  $B_{\mathbf{e}_i}$  equals to 0, which is expressed by  $\mathcal{S}_3$ .

Consider now any matrix  $B_{\mathbf{e}_i}$ . Its size is  $n \times d$ , thus any row of  $B_{\mathbf{e}_i}$  can be written as an element of  $\mathbb{F}_{2^d}$ . Let us define the matrix  $\mathfrak{B}_{\circ} = [B_{\mathbf{e}_1} \dots B_{\mathbf{e}_n}] \in (\mathbb{F}_{2^d})^{n \times n}$ . From Condition i), ii) and iii), above, we have the following properties for  $\mathfrak{B}_{\circ}$ :

- (I)  $\mathfrak{B}_{\circ}$  is full rank over  $\mathbb{F}_2$ . This guarantees that  $\dim(U(T_{\circ})) = d$ .
- (II)  $\mathfrak{B}_{\circ}$  is symmetric. Being the  $i$ th row of  $B_{\mathbf{e}_j}$  equal to the  $j$ th row of  $B_{\mathbf{e}_i}$  we have that the entry  $(i, j)$  of  $\mathfrak{B}_{\circ}$  is equal to the entry  $(j, i)$ .
- (III)  $\mathfrak{B}_{\circ}$  is zero-diagonal, as the  $i$ th row of  $B_{\mathbf{e}_i}$  is zero for all  $i$ .

**Example 3.1.** Let  $N = 5$  and  $d = 2$  and consider the operation  $\circ$  defined by the following  $B_{\mathbf{e}_i}$ 's:

$$B_{\mathbf{e}_1} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}, B_{\mathbf{e}_2} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}, B_{\mathbf{e}_3} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, B_{\mathbf{e}_4} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, B_{\mathbf{e}_5} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Since we have  $n = N - d = 3$ , we just need to focus on  $B_{\mathbf{e}_1}$ ,  $B_{\mathbf{e}_2}$  and  $B_{\mathbf{e}_3}$ .

Representing the rows of the matrices as elements of  $\mathbb{F}_{2^2} = \{0, 1, \alpha, \alpha + 1\}$ , where  $\alpha^2 = \alpha + 1$ , we can rewrite the matrices as

$$B_{\mathbf{e}_1} = \begin{bmatrix} 0 \\ \alpha + 1 \\ \alpha + 1 \end{bmatrix}, B_{\mathbf{e}_2} = \begin{bmatrix} \alpha + 1 \\ 0 \\ \alpha \end{bmatrix}, B_{\mathbf{e}_3} = \begin{bmatrix} \alpha + 1 \\ \alpha \\ 0 \end{bmatrix}$$

and so we can rewrite  $\mathfrak{B}_{\circ}$  as

$$\mathfrak{B}_{\circ} = [B_{\mathbf{e}_1} \ B_{\mathbf{e}_2} \ B_{\mathbf{e}_3}] = \begin{bmatrix} 0 & \alpha + 1 & \alpha + 1 \\ \alpha + 1 & 0 & \alpha \\ \alpha + 1 & \alpha & 0 \end{bmatrix}$$

In Theorem 2.8 we have a one-to-one correspondence between the matrices  $B_{\mathbf{e}_i}$ 's and the points of  $\mathcal{V}(\mathcal{I}_{n,d})$ . Thanks to the  $\mathfrak{B}_{\circ}$  construction, we have a one-to-one correspondence between the matrices  $\mathfrak{B}_{\circ} \in (\mathbb{F}_{2^d})^{n \times n}$  having the

above characteristics (I), (II), (III) and the groups  $T_\circ$  with  $U(T_\circ)$  generated by  $\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}$ . That is, defining

$$\mathcal{M}_{n,d} = \{\mathfrak{B}_\circ \in (\mathbb{F}_{2^d})^{n \times n} \mid \mathfrak{B}_\circ \text{ satisfies conditions (I), (II), and (III)}\}$$

we have  $|\mathcal{M}_{n,d}| = |\mathcal{V}(\mathcal{I}_{n,d})|$ . Thus, we aim to estimate  $|\mathcal{M}_{n,d}|$ .

We recall the following result given in [10].

**Proposition 3.2.** *Let  $q$  a power of 2. The number of  $n \times n$  symmetric invertible matrices over  $\mathbb{F}_q$  with zero diagonal is*

$$\begin{cases} q^{\binom{n}{2}} \prod_{j=1}^{\lceil \frac{n-1}{2} \rceil} (1 - q^{1-2j}) & n \text{ even} \\ 0 & n \text{ odd} \end{cases}.$$

**Proposition 3.3.** *Let  $N = n + d$  and  $V = (\mathbb{F}_2)^N$ , with  $n \geq 2$  and  $d \geq 1$ . Then we can define a lower bound  $\nu(n, d) \leq |\mathcal{M}_{n,d}|$  by*

$$\nu(n, d) = \begin{cases} q^{\binom{n}{2}} \prod_{j=1}^{\lceil \frac{n-1}{2} \rceil} (1 - q^{1-2j}) & \text{with } q = 2^d, n \text{ even} \\ (q^{n-1} - 2^{n-1}) q^{\binom{n-1}{2}} \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j}) & \text{with } q = 2^d, n \text{ odd} \end{cases}$$

*Proof.* We require that  $\mathfrak{B}_\circ$  is full rank over  $\mathbb{F}_2$ , thus if it is invertible this condition is satisfied. Then the value given in Proposition 3.2 is a lower bound on the number of all acceptable matrices  $\mathfrak{B}_\circ$ . However, if  $n$  is odd from Proposition 3.2 we would have only 0 as lower bound.

To tackle the  $n$ -odd case, we want to reduce it to the  $n$ -even case. We show how it is possible to construct a matrix  $\mathfrak{B}_\circ$  for fixed values  $n$  and  $d$  starting from a given  $\mathfrak{B}'_\circ$  defined for values  $n' = n - 1$  and  $d' = d$ . Indeed, let  $\mathfrak{B}'_\circ \in (\mathbb{F}_{2^d})^{n-1 \times n-1}$  be such that it is full rank over  $\mathbb{F}_2$ , symmetric and zero-diagonal. We need to construct the first row (the first column is the transpose of this) of  $\mathfrak{B}_\circ$ , setting all the others equal to the rows of  $\mathfrak{B}'_\circ$ . That is,

$$\mathfrak{B}_\circ = \begin{bmatrix} 0 & \mathbf{b} \\ \mathbf{b}^\top & \mathfrak{B}'_\circ \end{bmatrix}.$$

Then, we need to verify how many possible  $\mathfrak{B}_\circ$  we can construct starting from  $\mathfrak{B}'_\circ$ , or equivalently how many vectors  $\mathbf{b} \in (\mathbb{F}_{2^d})^{n-1}$  we can use, in order to enforce  $\mathfrak{B}_\circ \in \mathcal{M}_{n,d}$ .

As  $\mathfrak{B}'_\circ$  is full rank over  $\mathbb{F}_2$ , summing the rows of this matrix we can create  $2^{n-1} - 1$  non-zero vectors. Thus, we can choose  $2^{d(n-1)} - 2^{n-1}$  different vectors  $\mathbf{b}$  of  $(\mathbb{F}_{2^d})^{n-1}$  to construct  $\mathfrak{B}_\circ$ . Hence, for any  $n$  we have  $(2^{d(n-1)} - 2^{n-1})|\mathcal{M}_{n-1,d}| \leq |\mathcal{M}_{n,d}|$  and, in particular, for  $n$  odd we have at least  $(2^{d(n-1)} - 2^{n-1})\nu(n-1, d)$  possible matrices  $\mathfrak{B}_\circ$ .  $\square$

**Proposition 3.4.** *Let  $N = n + d$  and  $V = (\mathbb{F}_2)^N$ . Then*

$$|\mathcal{M}(\mathcal{I}_{n,d})| = \begin{cases} 2^d - 1 & n = 2 \\ (2^d + 3)(2^d - 1)(2^d - 2) & n = 3 \\ 2^{\binom{n}{2}} \prod_{j=1}^{\lceil \frac{n-1}{2} \rceil} (1 - 2^{1-2j}) & d = 1 \text{ and } n \text{ even} \\ 0 & d = 1 \text{ and } n \text{ odd} \end{cases}$$

*Proof.* The cases  $n = 2$  and  $n = 3$  were already proved in [1]. We restate here the proof using symmetric matrices, obtaining a shorter and more clear proof.

If  $n = 2$  then

$$\mathfrak{B}_\circ = \begin{bmatrix} 0 & \mathbf{x} \\ \mathbf{x} & 0 \end{bmatrix}$$

with  $\mathbf{x} \in \mathbb{F}_{2^d}$ . Of course the only vector that we have to avoid is the zero vector. Thus we can use  $2^d - 1$  vectors  $\mathbf{x}$  to construct  $\mathfrak{B}_\circ$ .

If  $n = 3$  then

$$\mathfrak{B}_\circ = \begin{bmatrix} 0 & \mathbf{y} & \mathbf{z} \\ \mathbf{y} & 0 & \mathbf{x} \\ \mathbf{z} & \mathbf{x} & 0 \end{bmatrix}$$

$\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{F}_{2^d}$  such that  $\mathfrak{B}_\circ \in \mathcal{M}_{n,d}$ . Let us consider the different cases:

- $\mathbf{x} \neq 0$  :  $2^d - 1$  possible values for  $\mathbf{x}$ ,
  - $\mathbf{y} = 0$  : so  $\mathbf{z} \notin \{0, \mathbf{x}\}$  and so we get  $2^d - 2$  possible values
  - $\mathbf{y} = \mathbf{x}$  : so  $\mathbf{z} \notin \{0, \mathbf{x}\}$  and so we get  $2^d - 2$  possible values
  - $\mathbf{y} \neq \mathbf{x}$  : so  $\mathbf{z}$  can be any element,  $\mathbf{y} \notin \{0, \mathbf{x}\}$ . So  $2^d(2^d - 2)$  possible pairs  $(\mathbf{y}, \mathbf{z})$ .
- $\mathbf{x} = 0$  : we have  $\mathbf{y} \neq 0$  and  $\mathbf{z} \notin \{0, \mathbf{y}\}$  and so we get  $(2^d - 1)(2^d - 2)$  possible pairs  $(\mathbf{y}, \mathbf{z})$ .

Summing all the possible triple  $(\mathbf{x}, \mathbf{y}, \mathbf{z})$ , we get

$$(2^d - 1)(2^d - 2 + 2^d - 2 + 2^d(2^d - 2)) + (2^d - 1)(2^d - 2) = (2^d - 1)(2^d - 2)(2^d + 3)$$

Now, let  $n \geq 2$  and fix  $d = 1$ . As we consider  $\mathbb{F}_{2^d} = \mathbb{F}_2$ , we have that  $\mathcal{M}_{n,d}$  is the set of symmetric matrices having all zeros on the diagonal and invertible. From Proposition 3.2 we have our claim.  $\square$

Let us compare the upper bound and the lower bound on the cardinality of  $\mathcal{M}_{n,d}$ . We recall a theorem on the infinite product convergence criteria, corollary of the monotone convergence theorem (more details can be found in [11]).

**Lemma 3.5.**

Let  $\{a_j\}_{j \in \mathbb{N}} \subseteq \mathbb{R}_+$ . Then  $\prod_{j=1}^{\infty} a_j$  converges if and only if  $\sum_{j=1}^{\infty} \ln(a_j)$  converges. Moreover, if  $a_j \geq 1$ , and let  $a_j = 1 + p_j$ . Then

$$1 + \sum_{j=1}^{\infty} p_j \leq \prod_{j=1}^{\infty} (1 + p_j) \leq e^{\sum_{j=1}^{\infty} p_j},$$

that is, the infinite product converges if and only if the infinite sum of the  $p_j$  converges.

**Proposition 3.6.** Let  $\mu(n, d)$  be the bound given in Proposition 2.9 and let  $\nu(n, d)$  be the bound given in Proposition 3.3. Let  $q = 2^d$  and  $d \geq 2$ . If  $n$  is even, then

$$\frac{\mu(n, d)}{\nu(n, d)} \leq \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \leq e^{\frac{q+1}{q(q-1)}}$$

if  $n$  is odd, then

$$\frac{\mu(n, d)}{\nu(n, d)} \leq 2 \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \leq 2e^{\frac{q+1}{q(q-1)}}.$$

*Proof.* We have

$$\begin{aligned} \mu(n, d) &= q^{\binom{n}{2}} - 1 - \sum_{r=1}^{n-2} \binom{n}{r} (q-1)^{\binom{n-r}{2}} \\ &\leq q^{\binom{n}{2}}. \end{aligned}$$

Consider the case  $n$  even

$$\begin{aligned} \frac{\mu(n, d)}{\nu(n, d)} &\leq \frac{q^{\binom{n}{2}}}{q^{\binom{n}{2}} \prod_{j=1}^{\lceil \frac{n-1}{2} \rceil} (1 - q^{1-2j})} \\ &= \prod_{j=1}^{\lceil \frac{n-1}{2} \rceil} \frac{1}{(1 - q^{1-2j})} \\ &\leq \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})}, \end{aligned}$$

since we can write

$$\frac{1}{1 - q^{1-2j}} = \frac{q^{2j-1}}{q^{2j-1} - 1} = 1 + \frac{q}{q^{2j} - q}$$

. Defining

$$p_j := \frac{q}{q^{2j} - q}$$

we trivially have that  $p_n \geq 0$  and

$$1 + \sum_{j=1}^{\infty} p_j \leq \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \leq e^{\sum_{j=1}^{\infty} p_j}.$$

We have for any  $j \geq 2$

$$p_j \leq \frac{1}{q^j}.$$

From this, we get

$$\sum_{i=1}^{\infty} p_i \leq p_1 + \sum_{i=2}^{\infty} \frac{1}{q^i} = \frac{2}{q-1} - \frac{1}{q} = \frac{q+1}{q(q-1)}$$

and so

$$\prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \leq e^{\frac{q+1}{q(q-1)}}.$$

Now let  $n$  be odd,

$$\frac{\mu(n, d)}{\nu(n, d)} \leq \frac{q^{\binom{n}{2}}}{(q^{n-1} - 2^{n-1}) \cdot q^{\binom{n-1}{2}} \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})}.$$

Being  $\binom{j}{i} = \binom{j-1}{i-1} + \binom{j-1}{i}$ ,  $n \geq 2$  and  $d \geq 2$  we have

$$\begin{aligned} \frac{\mu(n, d)}{\nu(n, d)} &\leq \frac{q^{n-1}}{(q^{n-1} - 2^{n-1}) \cdot \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})} \\ &= \frac{1}{\left(1 - \frac{2^{n-1}}{q^{n-1}}\right) \cdot \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})} \\ &\leq \frac{1}{\left(1 - \frac{2^{n-1}}{q^{n-1}}\right)} \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \\ &\leq 2 \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})} \leq 2e^{\frac{q+1}{q(q-1)}}. \end{aligned}$$

□

Note that in Proposition 3.6, the comparison for the case  $d = 1$  is avoided as the lower bound is the exact value of  $|\mathcal{M}_{n,d}|$ .

**Corollary 3.7.** *For all possible values of  $n \geq 2$  and  $d \geq 2$  we have*

$$\lim_{n \rightarrow \infty} \frac{\mu(n, d)}{\nu(n, d)} = \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})}$$

and

$$\lim_{d \rightarrow \infty} \frac{\mu(n, d)}{\nu(n, d)} = 1.$$

*Proof.* Consider the case  $n$  odd, then

$$\begin{aligned}
\frac{\mu(n, d)}{\nu(n, d)} &= \frac{q^{\binom{n}{2}} \left( 1 - \frac{1}{q^{\binom{n}{2}}} - \frac{\sum_{r=1}^{n-2} \binom{n}{r} \prod_{i=1}^{\binom{n-r}{2}} (q-1)}{q^{\binom{n}{2}}} \right)}{(q^{n-1} - 2^{n-1}) \cdot q^{\binom{n-1}{2}} \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})} \\
&= \frac{\left( 1 - \frac{1}{q^{\binom{n}{2}}} - \frac{\sum_{r=1}^{n-2} \binom{n}{r} \prod_{i=1}^{\binom{n-r}{2}} (q-1)}{q^{\binom{n}{2}}} \right)}{\left( 1 - \frac{2^{n-1}}{q^{n-1}} \right) \cdot \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})} \\
&= \frac{\left( 1 - \frac{1}{q^{\binom{n}{2}}} - \frac{\sum_{r=1}^{n-2} \binom{n}{r} \prod_{i=1}^{\binom{n-r}{2}} (q-1)}{q^{\binom{n}{2}}} \right)}{\left( 1 - \frac{2^{n-1}}{q^{n-1}} \right)} \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} \frac{1}{(1 - q^{1-2j})}
\end{aligned}$$

Consider

$$\begin{aligned}
\frac{\sum_{r=1}^{n-2} \binom{n}{r} \prod_{i=1}^{\binom{n-r}{2}} (q-1)}{q^{\binom{n}{2}}} &= \frac{\sum_{r=1}^{n-2} \binom{n}{r} (q-1)^{\binom{n-r}{2}}}{q^{\binom{n}{2}}} \\
&\leq \sum_{r=1}^{n-2} \binom{n}{r} q^{\binom{n-1}{2} - \binom{n}{2}} \\
&\leq \frac{2^{n-2}}{q^{n-1}}.
\end{aligned} \tag{3}$$

This implies that the limit of (3), as  $n$  approaches infinity, is 0 and then the limit

$$\lim_{n \rightarrow \infty} \frac{q^{\binom{n}{2}} \left( 1 - \frac{1}{q^{\binom{n}{2}}} - \frac{\sum_{r=1}^{n-2} \binom{n}{r} \prod_{i=1}^{\binom{n-r}{2}} (q-1)}{q^{\binom{n}{2}}} \right)}{(q^{n-1} - 2^{n-1}) \cdot q^{\binom{n-1}{2}} \prod_{j=1}^{\lceil \frac{n-2}{2} \rceil} (1 - q^{1-2j})} = \prod_{j=1}^{\infty} \frac{1}{(1 - q^{1-2j})}.$$

The case  $n$  even is similar.

Moreover, from the proof of Proposition 3.6 we have

$$1 \leq \frac{\mu(n, d)}{\nu(n, d)} \leq \begin{cases} e^{\frac{q+1}{q(q-1)}} & \text{if } n \text{ even} \\ \frac{1}{\left(1 - \frac{2^{n-1}}{q^{n-1}}\right)} e^{\frac{q+1}{q(q-1)}} & \text{if } n \text{ odd} \end{cases}$$

and immediately we have that

$$\lim_{d \rightarrow \infty} \frac{\mu(n, d)}{\nu(n, d)} = 1.$$

□

#### 4. On hidden sums for linear maps

In this section we investigate Problem 3 given in Section 2 page 4. In particular we want to see if, for a given  $\lambda \in \text{GL}(V, +)$ , it is possible to individuate an alternative sum  $\circ$  such that  $\lambda \in \text{GL}(V, \circ)$ .

**Proposition 4.1.** *Let  $T_\circ \subseteq \text{AGL}(V, +)$  and  $\lambda \in \text{GL}(V, +) \cap \text{GL}(V, \circ)$  then  $U(T_\circ)$  is invariant under the action of  $\lambda$ , i.e.  $U(T_\circ)\lambda = U(T_\circ)$ .*

*Proof.*  $\lambda$  is linear with respect to both  $+$  and  $\circ$ , thus let  $\mathbf{y} \in U(T_\circ)$  for all  $\mathbf{x}$  we have

$$\mathbf{x} \circ \mathbf{y} \lambda = \mathbf{x} \lambda^{-1} \lambda \circ \mathbf{y} \lambda = (\mathbf{x} \lambda^{-1} \circ \mathbf{y}) \lambda = (\mathbf{x} \lambda^{-1} + \mathbf{y}) \lambda = \mathbf{x} + \mathbf{y} \lambda.$$

That implies  $\mathbf{y} \lambda \in U(T_\circ)$ , and so  $U(T_\circ)\lambda \subseteq U(T_\circ)$ .  $\square$

**Proposition 4.2.** *Let  $T_\circ \subseteq \text{AGL}(V, +)$  and  $\lambda \in \text{GL}(V, +)$ . Then  $\lambda$  is in  $\text{GL}(V, +) \cap \text{GL}(V, \circ)$  if and only if for all  $\mathbf{x} \in V$  we have*

$$\kappa_{\mathbf{x}} \lambda = \lambda \kappa_{\mathbf{x}} \lambda, \quad (4)$$

where  $\tau_{\mathbf{x}} = \kappa_{\mathbf{x}} \sigma_{\mathbf{x}}$ .

*Proof.* Let  $\mathbf{y}$  be fixed, then for all  $\mathbf{x}$  we have

$$(\mathbf{x} \circ \mathbf{y}) \lambda = \mathbf{x} \lambda \circ \mathbf{y} \lambda$$

where

$$\begin{aligned} (\mathbf{x} \circ \mathbf{y}) \lambda &= (\mathbf{x} \kappa_{\mathbf{y}} + \mathbf{y}) \lambda \\ &= \mathbf{x} \kappa_{\mathbf{y}} \lambda + \mathbf{y} \lambda \end{aligned}$$

and

$$\mathbf{x} \lambda \circ \mathbf{y} \lambda = \mathbf{x} \lambda \kappa_{\mathbf{y} \lambda} + \mathbf{y} \lambda$$

Imposing the equality we get

$$\mathbf{x}(\kappa_{\mathbf{y}} \lambda) = \mathbf{x} \lambda \kappa_{\mathbf{y} \lambda}.$$

Vice versa let  $\mathbf{x}, \mathbf{y} \in V$

$$\begin{aligned} (\mathbf{x} \circ \mathbf{y}) \lambda &= \mathbf{x} \kappa_{\mathbf{y}} \lambda + \mathbf{y} \lambda \\ &= \mathbf{x} \lambda \kappa_{\mathbf{y} \lambda} + \mathbf{y} \lambda \\ &= \mathbf{x} \lambda \circ \mathbf{y} \lambda. \end{aligned}$$

$\square$

Now we will characterize the linear maps that are also linear for an operation  $\circ$  such that  $U(T_\circ)$  is generated by the last elements of the canonical basis.

**Proposition 4.3.** *Let  $V = (\mathbb{F}_2)^N$ , with  $N = n + d$ ,  $n \geq 2$  and  $d \geq 1$ . Let  $T_o \subseteq \text{AGL}(V, +)$  with  $U(T_o) = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$ . Let  $\lambda \in \text{GL}(V, +)$ . Then  $\lambda \in \text{GL}(V, +) \cap \text{GL}(V, \circ)$  if and only if*

$$\lambda = \begin{bmatrix} \Lambda_1 & \Lambda_2 \\ 0 & \Lambda_3 \end{bmatrix},$$

with  $\Lambda_1 \in \text{GL}((\mathbb{F}_2)^n)$ ,  $\Lambda_3 \in \text{GL}((\mathbb{F}_2)^d)$ ,  $\Lambda_2$  any matrix and for all  $\mathbf{x} \in V$   $B_{\mathbf{x}}\Lambda_3 = \Lambda_1 B_{\mathbf{x}\lambda}$  (see Theorem 2.6 for the notation of  $B_{\mathbf{x}}$ ).

*Proof.* Let

$$\lambda = \begin{bmatrix} \Lambda_1 & \Lambda_2 \\ \Lambda_4 & \Lambda_3 \end{bmatrix},$$

$\Lambda_1 \in (\mathbb{F}_2)^{n \times n}$ ,  $\Lambda_3 \in (\mathbb{F}_2)^{d \times d}$ ,  $\Lambda_2 \in (\mathbb{F}_2)^{n \times d}$  and  $\Lambda_4 \in (\mathbb{F}_2)^{d \times n}$

From Proposition 4.1 we have that  $\Lambda_4 = 0$  as  $U(T_o)$  is  $\lambda$ -invariant. Thus, being  $\lambda$  invertible  $\Lambda_1 \in \text{GL}((\mathbb{F}_2)^n)$  and  $\Lambda_3 \in \text{GL}((\mathbb{F}_2)^d)$ .

By standard matrix multiplication it can be verified that (4) and  $B_{\mathbf{x}}\Lambda_3 = \Lambda_1 B_{\mathbf{x}\lambda}$  are equivalent. Then, we need to show that this condition does not depends on the matrix  $\Lambda_2$ . Indeed, let  $\mathbf{x} = (x_1, \dots, x_n, x_{n+1}, \dots, x_{n+d}) \in V$ , and define  $\bar{\mathbf{x}} = (x_1, \dots, x_n)$  and  $\mathbf{x}' = (x_{n+1}, \dots, x_{n+d})$ , i.e.  $\mathbf{x} = (\bar{\mathbf{x}}, \mathbf{x}')$ . As reported in Section 3 page 6, we have  $\kappa_{\mathbf{x}} = \kappa_{(\bar{\mathbf{x}}, 0)}$ , which implies  $B_{\mathbf{x}} = B_{(\bar{\mathbf{x}}, 0)}$ , for all  $\mathbf{x} \in V$ . Then,  $B_{\mathbf{x}}\Lambda_3 = \Lambda_1 B_{\mathbf{x}\lambda}$  for all  $\mathbf{x} \in V$  if and only if  $B_{(\bar{\mathbf{x}}, 0)}\Lambda_3 = \Lambda_1 B_{(\bar{\mathbf{x}}\Lambda_1, 0)}$  for all  $\bar{\mathbf{x}} = (x_1, \dots, x_n)$ . So we have

$$\begin{bmatrix} \Lambda_1 & \Lambda_2 \\ 0 & \Lambda_3 \end{bmatrix} \in \text{GL}(V, \circ),$$

for any  $\Lambda_2$ . □

In the following, as  $\Lambda_2$  in Proposition 4.3 could be any matrix, we will consider linear maps of type

$$\begin{bmatrix} \Lambda_1 & * \\ 0 & \Lambda_2 \end{bmatrix} \in \text{GL}(V, \circ),$$

with  $\Lambda_1 \in \text{GL}((\mathbb{F}_2)^n)$ ,  $\Lambda_2 \in \text{GL}((\mathbb{F}_2)^d)$  and  $*$  denotes for any matrix of size  $n \times d$ .

**Remark 4.4.** *From the propositions above, if we want to find an operation  $\circ$  that linearizes a linear map  $\lambda \in \text{GL}(V, +)$ , i.e. we want to enforce*

$$\begin{bmatrix} \Lambda_1 & * \\ 0 & \Lambda_2 \end{bmatrix} \in \text{GL}(V, \circ),$$

then we have to construct some matrices  $B_{\mathbf{x}}$  such that  $B_{\mathbf{x}}\Lambda_2 = \Lambda_1 B_{\mathbf{x}\lambda}$  for all  $\mathbf{x}$ . Moreover, as the standard vectors  $\mathbf{e}_i$ 's form a basis for the operation  $\circ$ ,

then we need to individuate only the matrices  $B_{\mathbf{e}_i}$ , so that  $B_{\mathbf{e}_i}\Lambda_2 = \Lambda_1 B_{\mathbf{e}_i}\lambda$ , and in particular that

$$B_{\mathbf{e}_i}\Lambda_2 = \Lambda_1 B_{\mathbf{e}_i}\lambda = \Lambda_1 \left( \sum_{i=1}^n c_i B_{\mathbf{e}_i} \right),$$

where  $c_1, \dots, c_n$  are the first components of the vector  $\mathbf{e}_i\lambda$ .

**Algorithm 4.1.**

INPUT:

$$\lambda = \begin{bmatrix} \Lambda_1 & * \\ 0 & \Lambda_2 \end{bmatrix},$$

with  $\Lambda_1 \in \text{GL}((\mathbb{F}_2)^n)$ ,  $\Lambda_2 \in \text{GL}((\mathbb{F}_2)^d)$

OUTPUT: all possible hidden sums such that:

- $T_{\circ} \subseteq \text{AGL}(V, +)$ ,  $T_+ \subseteq \text{AGL}(V, \circ)$ ,
- $U(T_{\circ})$  contains  $\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}$ ,
- $\lambda \in \text{GL}(V, \circ)$ .

ALGORITHM STEPS:

I) Consider the canonical basis  $\mathbf{e}_1, \dots, \mathbf{e}_{n+d}$  and compute  $\mathbf{e}_1\lambda, \dots, \mathbf{e}_n\lambda$

II) Solve the linear system given by the equations:

1. for all  $i = 1, \dots, n$

$$B_{\mathbf{e}_i}\Lambda_2 = \Lambda_1 B_{\mathbf{e}_i}\lambda = \Lambda_1 \left( \sum_{i=1}^n c_i B_{\mathbf{e}_i} \right),$$

(where  $c_1, \dots, c_n$  are the first components of the vector  $\mathbf{e}_i\lambda$ ).

2. for all  $i = n+1, \dots, n+d$

$$B_{\mathbf{e}_i} = 0,$$

3. for all  $i = 1, \dots, n$

$$\bar{\mathbf{e}}_i B_{\mathbf{e}_i} = 0,$$

(here  $\bar{\mathbf{e}}_i$  is the truncation of  $\mathbf{e}_i$  with respect to the first  $n$  coordinates)

4. for all  $i, j = 1, \dots, n$

$$\bar{\mathbf{e}}_i B_{\mathbf{e}_j} = \bar{\mathbf{e}}_j B_{\mathbf{e}_i},$$

III) return the solutions  $\{B_{\mathbf{e}_i}\}_{i=1, \dots, n+d}$ .

Note that, from Algorithm 4.1 we obtain operations  $\circ$  such that  $\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\} \subseteq U(T_{\circ})$ . Indeed, we required that for  $n+1 \leq i \leq n+d$ ,  $B_{\mathbf{e}_i} = 0$ , but we did not require that the combinations of  $B_{\mathbf{e}_1}, \dots, B_{\mathbf{e}_n}$  are non-zero. So, if we want to construct hidden sums with  $\dim(U(T_{\circ})) = d$ , we just process the solution of Algorithm 4.1 and discharge the  $\mathfrak{B}_{\circ}$ 's such that  $\text{rank}_{\mathbb{F}_2}(\mathfrak{B}_{\circ}) < n$ .

#### 4.1. Complexity of our search algorithm

To solve the linear system in our algorithm, we represent the matrices as vectors:

$$\begin{bmatrix} b_{1,1} & \cdots & b_{1,d} \\ \vdots & \ddots & \vdots \\ b_{n,1} & \cdots & b_{n,d} \end{bmatrix} \longleftrightarrow \begin{bmatrix} \overbrace{b_{1,1}, \dots, b_{1,d}}^{\text{1-st row}} & \overbrace{b_{2,1}, \dots, b_{2,d}}^{\text{2-nd row}} & \cdots & \overbrace{b_{n,1}, \dots, b_{n,d}}^{\text{n-th row}} \end{bmatrix}$$

- From  $i = 1, \dots, n$ ,  $\sum_j c_j \Lambda_1 B_{\mathbf{e}_j} = B_{\mathbf{e}_i} \Lambda_2$  we have  $n^2 d$  linear equation in  $n^2 d$  variables.
- From  $\bar{\mathbf{e}}_i B_{\mathbf{e}_i} = 0$  and  $\bar{\mathbf{e}}_i B_{\mathbf{e}_j} = \bar{\mathbf{e}}_j B_{\mathbf{e}_i}$  we obtain  $\binom{n+1}{2} d$  linear equations, in the same variables.

So, we only need to find a solution of a binary linear system of size

$$\left( n^2 d + \binom{n+1}{2} d \right) \cdot n^2 d.$$

Therefore, we have immediately the following result

**Proposition 4.5.** *The time complexity of Algorithm 4.1 is  $\mathcal{O}(n^6 d^3)$  and the space complexity is  $\mathcal{O}(l \cdot 2^{d-1} n^2)$  where  $l$  is the dimension of the solution subspace.*

In Table 1 we report some timings for different dimensions of the message space  $V$ , fixing the value of  $d$  equal to 2.

| Dimension of $V$ | $n, d$ | Timing in second |
|------------------|--------|------------------|
| 64               | 62, 2  | 32.620 seconds   |
| 80               | 78, 2  | 84.380 seconds   |
| 96               | 94, 2  | 188.200 seconds  |
| 112              | 110, 2 | 338.590 seconds  |
| 128              | 126, 2 | 616.670 seconds  |

Table 1: Computation timing for a Mac Book Pro 15" early 2011, 4 GB Ram, Intel i7 2.00 Ghz.

## 5. Hidden sum for PRESENT's mixing layer

Here we report our results on the search for a hidden sum suitable for the mixing layer of PRESENT,  $\lambda_P$ , which is defined by the permutation reported in Table 2.

|              |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |    |
|--------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| $i$          | 1  | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9  | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| $i\lambda_P$ | 1  | 17 | 33 | 49 | 1  | 18 | 34 | 50 | 3  | 19 | 35 | 51 | 4  | 20 | 36 | 52 |
| $i$          | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 |
| $i\lambda_P$ | 5  | 21 | 37 | 53 | 6  | 22 | 38 | 54 | 7  | 23 | 39 | 55 | 8  | 24 | 40 | 56 |
| $i$          | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 |
| $i\lambda_P$ | 9  | 25 | 41 | 57 | 10 | 26 | 42 | 58 | 11 | 27 | 43 | 59 | 12 | 28 | 44 | 60 |
| $i$          | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 | 61 | 62 | 63 | 64 |
| $i\lambda_P$ | 13 | 29 | 45 | 61 | 14 | 30 | 46 | 62 | 15 | 31 | 47 | 63 | 16 | 32 | 48 | 64 |

Table 2: PRESENT mixing layer

Algorithm 4.1 requires in input a linear function in the block form:

$$\lambda' = \begin{pmatrix} \Lambda_1 & * \\ 0 & \Lambda_2 \end{pmatrix} \quad (5)$$

with  $\Lambda_1 \in \text{GL}((\mathbb{F}_2)^n)$  and  $\Lambda_2 \in \text{GL}((\mathbb{F}_2)^d)$  for some integers  $n$  and  $d$ .

Note that a matrix as in (5) is such that the space  $U' = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$  is  $\lambda'$ -invariant. So, in order to transform a mixing layer  $\lambda$  into one as in (5), it is necessary to individuate a subspace  $U$  such that  $U\lambda = U$ . Then, by conjugating  $\lambda$  by a linear map  $\pi$  such that  $U\pi = \text{Span}\{\mathbf{e}_{n+1}, \dots, \mathbf{e}_{n+d}\}$ , we will have  $\pi\lambda\pi^{-1}$  as in (5). From a group  $T_\circ$  obtained from Algorithm 4.1 for the map  $\pi\lambda\pi^{-1}$  we will obtain a hidden sum for  $\lambda$ , that is,  $\pi^{-1}T_\circ\pi$ .

For this reason we consider the matrix given by the permutation

$$\pi_P = (1, 61)(22, 62)(43, 63) \in \text{Sym}(\{i \mid i = 1, \dots, 64\})$$

so that

$$\pi_P \lambda_P \pi_P^{-1} = \begin{pmatrix} \Lambda_1 & 0 \\ 0 & I_4 \end{pmatrix} = \hat{\lambda}_P$$

We can now apply Algorithm 4.1 and obtain all the possible operations that linearize  $\hat{\lambda}_P$ . The operation space will be denoted with  $O$ .

The time required to compute the operation space  $O$  is  $\sim 10.420$  seconds and it is generated by 2360 60-tuples of  $60 \times 4$  matrices. So the number of operations that linearize  $\hat{\lambda}_P$ , in the form described in Theorem 2.6, is  $|O| = 2^{2360}$ .

We take a random operation  $\circ$  (Table 3) obtained by our algorithm. The operation has  $\text{rank}_{\mathbb{F}_2}(\mathfrak{B}_\circ) = 60 = n$  and so the operation is such that  $\dim(U(T_\circ)) = 4 = d$ . To compress the table, we represent every row of  $B_{e_i}$  as a number in  $\{0, \dots, (2^4 - 1)\}$ , and, as we did in Remark 4.1, row  $i$  represents the matrix  $B_{\mathbf{e}_i}$ .



## 6. Conclusions

Continuing the study of [1], here we focused on the class of hidden sums such that  $T_{\circ} \subseteq \text{AGL}(V, +)$  and  $T_{+} \subseteq \text{AGL}(V, \circ)$ , as these could be used to exploit new attacks on block ciphers.

We gave a lower bound on the number of these hidden sums, comparing this values with that given in [1].

In the second part, we dealt with the problem of individuating possible hidden sums for a given linear map, providing Algorithm 4.1 and showing an example on the case of the PRESENT's mixing layer.

A dual approach is under investigation in [9], where the authors take into consideration some S-boxes, which would be strong according to classical cryptanalysis, and identify some hidden sums that weaken significantly the non-linearity properties of the S-Boxes. Once the candidate hidden sums have been found, some (linear) mixing layers are constructed such that they are linear with respect to a hidden sum and the resulting cipher turns out to be attackable by the knowingly attacker.

## References

- [1] M. Calderini, M. Sala, Elementary abelian regular subgroups as hidden sums for cryptographic trapdoors, arXiv preprint arXiv:1702.00581.
- [2] A. Bogdanov, L. R. Knudsen, G. Leander, C. Paar, A. Poschmann, M. Robshaw, Y. Seurin, C. Viskelson, PRESENT: An Ultra-Lightweight Block Cipher, in: Proc. of CHES 2007, Vol. 7427 of LNCS, Springer, 2007, pp. 450–466.
- [3] D. Stinson, Cryptography, Theory and Practice, CRC Press, 1995.
- [4] J. Daemen, V. Rijmen, The design of Rijndael, Information Security and Cryptography, Springer-Verlag, Berlin, 2002, AES - the Advanced Encryption Standard.
- [5] A. Caranti, F. Dalla Volta, M. Sala, On some block ciphers and imprimitive groups, AAECC 20 (5-6) (2009) 229–350.
- [6] J. Daemen, V. Rijmen, The design of Rijndael: AES-the advanced encryption standard, Springer Science & Business Media, 2002.
- [7] R. Anderson, E. Biham, L. Knudsen, SERPENT: A New Block Cipher Proposal, in: Fast Software Encryption, Vol. 1372 of LNCS, Springer, 1998, pp. 222–238.
- [8] K. G. Paterson, Imprimitive permutation groups and trapdoors in iterated block ciphers, in: Fast software encryption, Vol. 1636 of LNCS, Springer, Berlin, 1999, pp. 201–214.
- [9] R. Civino, in preparation, Ph.D. thesis, University of TRENTO (2016).
- [10] J. MacWilliams, Orthogonal matrices over finite fields, The American Mathematical Monthly 76 (2) (1969) 152–164.  
URL <http://www.jstor.org/stable/2317262>
- [11] J. Stewart, Calculus, Cengage Learning, 2015.